



LIST OF REFERENCES CITED BY APPLICANT

(Use several sheets if necessary)

ATTY DOCKET NO.

10165-027-999

APPLICATION NO.

10/676,694

APPLICANT

Brines et al.

FILING DATE

September 30, 2003

GROUP

+647 1646

U.S. PATENT DOCUMENTS

*EXAMINER INITIAL		DOCUMENT NUMBER	DATE	NAME	CLASS	SUBCLASS	FILING DATE IF APPROPRIATE
/R.L./	A01	2004/0214236	10/28/04	Brines et al.			
	A02	2004/0122216	6/24/04	Nielsen et al.			
	A03	2003/0134798	7/17/03	Brines et al.			
	A04	2003/0104988	6/5/03	Brines et al.			
	A05	2003/0072737	4/17/03	Brines et al.			
	A06	2002/0031806	3/14/02	Lee			
	A07	6,531,121	3/11/03	Brines et al.			
	A08	5,763,198	6/9/98	Hirth et al.			
/R.L./	A09	5,292,654	3/8/94	Yoshimura et al.			

FOREIGN PATENT DOCUMENTS

		DOCUMENT NUMBER	DATE	COUNTRY	CLASS	SUBCLASS	TRANSLATION
							YES NO
/R.L./	B01	WO 04/096148	11/11/04	PCT			
/R.L./	B02	WO 02/053580	7/11/02	PCT			

OTHER REFERENCES (Including Author, Title, Date, Pertinent Pages, Etc.)

/R.L./	C01	Barber et al., 1994, "Erythropoietin and interleukin-2 activate distinct JAK kinase family members," Mol. Cell. Biol. 14(10):6506-6514
	C02	Barbone et al., 1997, "Mutagenesis studies of the human erythropoietin receptor. Establishment of structure-function relationships," J. Biol. Chem. 272(8):4985-4992
	C03	Bazan, 1989, "A novel family of growth factor receptors: a common binding domain in the growth hormone, prolactin, erythropoietin and IL-6 receptors, and the p75 IL-2 receptor beta-chain," Biochem. Biophys. Res. Commun. 164(2):788-795
	C04	Benit et al., 1993, "The 'WS motif' common to v-mpl and members of the cytokine receptor superfamily is dispensable for myeloproliferative leukemia virus pathogenicity," Oncogene 8(3):787-790
	C05	Bonsi et al., 1997, "An erythroid and megakaryocytic common precursor cell line (B1647) expressing both c-mpl and erythropoietin receptor (Epo-R) proliferates and modifies globin chain synthesis in response to megakaryocyte growth and development factor (MGDF) but not to erythropoietin (Epo)," Br. J. Haematol. 98(3):549-559
	C06	Boudot et al., 1999, "Erythropoietin induces glycosylphosphatidylinositol hydrolysis. Possible involvement of phospholipase c-gamma(2)," J. Biol. Chem. 274(48):33966-33972
	C07	Brizzi et al., 1991, "Hematopoietic growth factor receptors," Int. J. Cell. Cloning 9(4):274-300
	C08	Caravella et al., 1996, "A partial model of the erythropoietin receptor complex," Proteins 24(3):394-401
	C09	D'Andrea et al., 2000, "A model for assembly and activation of the GM-CSF, IL-3 and IL-5 receptors: insights from activated mutants of the common beta subunit," Exp. Hematol. 28(3):231-243
	C10	D'Andrea et al., 1998, "Dysregulated hematopoiesis and a progressive neurological disorder induced by expression of an activated form of the human common beta chain in transgenic mice," J. Clin. Invest. 102(11):1951-1960
/R.L./	C11	D'Andrea and Zon, 1990, "Erythropoietin receptor. Subunit structure and activation," J. Clin. Invest. 86(3):681-687

/R.L./	C12	Frank, 2002, "Receptor dimerization in GH and erythropoietin action--it takes two to tango, but how?" <i>Endocrinology</i> 143(1):2-10
	C13	Fung et al., 1990, "The human interleukin-2 receptor: insights into subunit structure and growth signal transduction," <i>Semin. Immunol.</i> 2(2):119-128
	C14	Grotzinger, 2002, "Molecular mechanisms of cytokine receptor activation," <i>Biochim. Biophys. Acta.</i> 1592(3):215-223
	C15	Hanazono et al., 1995, "Erythropoietin induces tyrosine phosphorylation of the b chain of the GM-CSF receptor," <i>Biochem. Biophys. Res. Commun.</i> 208(3):1060-1066
	C16	Harris et al., 1992, "Ligand binding properties of the human erythropoietin receptor extracellular domain expressed in <i>Escherichia coli</i> ," <i>J. Biol. Chem.</i> 267(21):15205-15209
	C17	Hassan et al., 1995, "Characteristic biological features of human megakaryoblastic leukaemia cell lines," <i>Leuk. Res.</i> 19(9):589-594
	C18	Imada et al., 1992, "Interleukin-2 (IL-2) induces erythroid differentiation and tyrosine phosphorylation in ELM-F1 cells transfected with a human IL-2 receptor beta chain cDNA," <i>Biochem. Biophys. Res. Commun.</i> 188(1):352-357
	C19	Itoh et al., 1990, "Cloning of an interleukin-3 receptor gene: a member of a distinct receptor gene family," <i>Science</i> 247(4940):324-327
	C20	Jenkins et al., 1999, "A cell type-specific constitutive point mutant of the common beta-subunit of the human granulocyte-macrophage colony-stimulating factor (GM-CSF), interleukin (IL)-3, and IL-5 receptors requires the GM-CSF receptor alpha-subunit for activation," <i>J. Biol. Chem.</i> 274(13):8669-8677
	C21	Jones et al., 1990, "Human erythropoietin receptor: cloning, expression, and biologic characterization," <i>Blood</i> 76(1):31-35
	C22	Jubinsky et al., 1997, "The beta chain of the interleukin-3 receptor functionally associates with the erythropoietin receptor," <i>Blood</i> 90(5):1867-1873
	C23	Jubinsky et al., 1996, "The beta c component of the granulocyte-macrophage colony-stimulating factor (GM-CSF)/interleukin 3 (IL-3)/IL-5 receptor interacts with a hybrid GM-CSF/erythropoietin receptor to influence proliferation and beta-globin mRNA expression," <i>Mol. Med.</i> 2(6):766-773
	C24	Kiritto et al., 2002, "Identification of the human erythropoietin receptor region required for Stat1 and Stat3 activation," <i>Blood</i> 99(1):102-110
	C25	Lai et al., 1996, "The molecular role of the common gamma c subunit in signal transduction reveals functional asymmetry within multimeric cytokine receptor complexes," <i>Proc. Natl. Acad. Sci. USA</i> 93(1):231-235
	C26	Lewis et al., 2004, "Opposing effects of PI3 kinase pathway activation on human myeloid and erythroid progenitor cell proliferation and differentiation in vitro," <i>Exp. Hematol.</i> 32(1):36-44
	C27	Lewis et al., 1996, "Molecular characterization of the 7q deletion in myeloid disorders," <i>Br. J. Haematol.</i> 93(1):75-80
	C28	Liu et al., 1994, "Multiple cytokines stimulate the binding of a common 145-kilodalton protein to Shc at the Grb2 recognition site of Shc," <i>Mol. Cell. Biol.</i> 14(10):6926-6935
	C29	Livnah et al., 1999, "Crystallographic evidence for preformed dimers of erythropoietin receptor before ligand activation," <i>Science</i> 283(5404):987-990
	C30	McClure et al., 2001, "GM-CSF binding to its receptor induces oligomerisation of the common beta-subunit," <i>Cytokine</i> 13(4):240-243
	C31	Means et al., 1996, "Inhibition of human erythroid colony-forming units by interferons alpha and beta: differing mechanisms despite shared receptor," <i>Exp. Hematol.</i> 24(2):204-208
	C32	Murakami et al., 1991, "Critical cytoplasmic region of the interleukin 6 signal transducer gp130 is conserved in the cytokine receptor family," <i>Proc. Natl. Acad. Sci. USA</i> 88(24):11349-11353
	C33	Murray, 1996, <i>Harpers Biochemistry</i> 24th ed. pp.524-526. Appleton & Lange, Ltd.
	C34	Naranda et al., 2002, "Activation of erythropoietin receptor through a novel extracellular binding site," <i>Endocrinology</i> 143(6):2293-2302
	C35	Nathan, 1994, "Studies of hybrid hematopoietic growth factor receptors," <i>Stem Cells</i> 12 Suppl 1:27-33
	C36	Noguchi et al., 1991, "Cloning of the human erythropoietin receptor gene," <i>Blood</i> 78(10):2548-2556
	C37	Penny and Forget, 1991, "Genomic organization of the human erythropoietin receptor gene," <i>Genomics</i> 11(4):974-980
	C38	Remy et al., 1999, "Erythropoietin receptor activation by a ligand-induced conformation change," <i>Science</i> 283(5404):990-993
	C39	Scott et al., 2000, "Reassessment of interactions between hematopoietic receptors using common beta-chain and interleukin-3-specific receptor beta-chain-null cells: no evidence of functional interactions with receptors for erythropoietin, granulocyte colony-stimulating factor, or stem cell factor," <i>Blood</i> 96(4):1588-1590
	C40	Shikama et al., 1996, "A constitutively activated chimeric cytokine receptor confers factor-independent growth in hematopoietic cell lines," <i>Blood</i> 88(2):455-464
	C41	Tojo et al., 1987, "Identification of erythropoietin receptors on fetal liver erythroid cells," <i>Biochem. Biophys. Res. Commun.</i> 148(1):443-448
	C41	Williamson et al., 1993, "Protein and lipid kinase activation cascades in interleukin-2 receptor signalling," <i>Semin. Immunol.</i> 5(5):337-344
	C43	Winkelmann et al., 1990, "The gene for the human erythropoietin receptor: analysis of the coding sequence and assignment to chromosome 19p," <i>Blood</i> 76(1):24-30
/R.L./	C44	Yamamura et al., 1992, "Distinct downstream signaling mechanism between erythropoietin receptor and interleukin-2 receptor," <i>EMBO J.</i> 11(13):4909-4915

/R.L./	C45	Yet et al, 1993, "The extracytoplasmic domain of the erythropoietin receptor forms a monomeric complex with erythropoietin," Blood 82(6):1713-1719
	C46	Yoshimura et al., 1996, "Mouse oncostatin M: an immediate early gene induced by multiple cytokines through the JAK-STATS pathway," EMBO J. 15(5):1055-1063
↓	C47	Yoshimura <i>et al.</i> , 1996, "Physician Education: The Erythropoietin Receptor and Signal Transduction," Oncologist 1(5):337-339
↓	C48	Yoshimura et al., 1995, "A novel cytokine-inducible gene CIS encodes an SH2-containing protein that binds to tyrosine-phosphorylated interleukin 3 and erythropoietin receptors," EMBO J. 14(12):2816-2826
/R.L./	C49	Zhu et al., 2002, "Detecting and responding to hypoxia," Nephrol. Dial. Transplant. 17 Suppl 1:3-7

EXAMINER	/Ruixiang Li/	DATE CONSIDERED	04/03/2009
*EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.			

ALL REFERENCES CONSIDERED EXCEPT WHERE LINED THROUGH. /R.L./